

# Efficacy and Safety of Tralokinumab Plus Topical Corticosteroids in Patients with Severe Atopic Dermatitis and Prior History of Dupilumab Treatment: A Post Hoc Subgroup Analysis from ECZTRA 7 Trial

Jan Gutermuth<sup>1</sup>, Andrew Pink<sup>2</sup>, Margitta Worm<sup>3</sup>, Lise Soldbro<sup>4</sup>, Thomas Mark<sup>4</sup>, Joshua Corriveau<sup>5</sup>, Christian Bjerregård Øland<sup>4</sup>, Shannon Schneider<sup>5</sup>, Stephan Weidinger<sup>6</sup>

<sup>1</sup>Universitair Ziekenhuis Brussel and Vrije Universiteit Brussel, Brussels, Belgium; <sup>2</sup>Guy's and St. Thomas' NHS Foundation Trust, London, UK; <sup>3</sup>Charité – Universitätsmedizin Berlin, Berlin, Germany; <sup>4</sup>LEO Pharma A/S, Ballerup, Denmark; <sup>5</sup>LEO Pharma Inc., Madison, NJ, USA; <sup>6</sup>University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany

## Introduction

- AD is a chronic inflammatory disease characterized by eczematous skin lesions and multiple symptoms including pruritus, sleep disturbance, and depression<sup>1-4</sup>
- Tralokinumab is a high-affinity, fully human, monoclonal antibody designed to specifically neutralize interleukin-13, a key driver of the underlying inflammation in AD<sup>5-7</sup>
- The Phase 3 ECZTRA 7 trial (NCT03761537) met its primary endpoint of EASI-75 at Week 16, confirming tralokinumab plus topical corticosteroids (TCS) is superior to placebo plus TCS in treating severe atopic dermatitis (AD) in patients not adequately controlled by, or with contraindications to, oral cyclosporine A
- There can be inadequate disease control with currently available treatment options, and many patients with severe AD continue to experience high disease burden


## Objective

- To describe the efficacy and safety of tralokinumab in a subgroup of ECZTRA 7 patients with prior history of dupilumab treatment

## Methods

- ECZTRA 7 was a randomized, double-blinded, multicenter, placebo-controlled Phase 3 trial (Figure 1)
- Key inclusion criteria for ECZTRA 7:
  - Adult patients with AD for ≥1 year with inadequate response to topical or documented systemic medication in the past year
  - Disease not adequately controlled with, or patients with contraindications to, use of oral cyclosporine A
  - AD involvement of ≥10% body surface area
  - EASI ≥20 and IGA ≥3 at screening and at baseline
  - Worst daily pruritus numeric rating scale (NRS) average score of ≥4 during the week prior to baseline

Figure 1. ECZTRA 7 trial design.



AD, atopic dermatitis; EASI, Eczema Area and Severity Index; IGA, Investigator's Global Assessment; Q2W, every 2 weeks; TCS, topical corticosteroid; TCI, topical calcineurin inhibitor.

- Eligible patients were randomized 1:1 to subcutaneous tralokinumab 300 mg every 2 weeks with TCS as needed or placebo with TCS as needed for a treatment period of 26 weeks, following a 600-mg loading dose on Day 0
- For this analysis, prior history of dupilumab treatment was confirmed and further details were collected via queries before trial unblinding
- Dupilumab-experienced patients were defined as those with a confirmed history of dupilumab use prior to the trial
- Cochran-Mantel-Haenszel with treatment as only strata was used for analysis

## Results

### Patient characteristics

- Dupilumab-experienced (n=14) and dupilumab-naive (n=263) cohorts had comparable baseline characteristics, except that median (interquartile range, IQR) age was 51.5 (43.0, 57.0) years for dupilumab-experienced patients and 33.0 (25.0, 45.0) years for dupilumab-naive patients (Table 1)
- Median (IQR) EASI and percentage of patients with an IGA of 4 were 35.5 (24.8, 39.6) and 57.1% among dupilumab-experienced patients and 28.7 (22.4, 39.5) and 49.0% among dupilumab-naive patients, respectively
- Among dupilumab-experienced patients, baseline and clinical characteristics were similar between the tralokinumab + TCS as needed (n=6) and placebo + TCS as needed (n=8) groups
  - 50% of patients in each of these two groups discontinued dupilumab due to either lack of efficacy or safety concerns

Table 1. Baseline demographics and clinical characteristics for randomized patients in ECZTRA 7

	Dupilumab-Naive			Dupilumab-Experienced								
	All (N=263)			All (N=14)		Tralokinumab + TCS (n=6)		Placebo + TCS (n=8)				
	n	Median	IQR	n	Median	IQR	n	Median	IQR	n	Median	IQR
AD duration, years	262	26.0	18.0, 34.0	14	34.0	15.0, 44.0	6	17	15.0, 43.0	8	34.0	21.0, 47.5
Age, years	263	33.0	25.0, 45.0	14	51.5	43.0, 57.0	6	50.0	43.0, 56.0	8	51.5	42.0, 62.5
BSA, %	263	52.0	35.0, 70.0	14	56.5	34.0, 70.0	6	58.5	50.0, 72.0	8	54.5	33.5, 65.0
DLQI	257	16.0	11.0, 21.0	14	15.0	8.0, 18.0	6	11.0	7.0, 16.0	8	16.5	10.5, 21.5
EASI	261	28.7	22.4, 39.5	14	35.5	24.8, 39.6	6	37.3	29.0, 39.6	8	32.3	23.5, 38.9
SCORAD	261	68.9	61.5, 78.9	14	73.6	61.2, 77.0	6	72.6	58.0, 73.8	8	76.7	64.7, 82.2
Worst daily pruritus NRS (weekly average)	259	7.4	6.6, 8.3	14	6.7	5.4, 8.0	6	5.9	5.3, 7.6	8	7.4	6.2, 8.9
IGA 4, n (%)	263	129 (49.0)		14	8 (57.1)		6	5 (83.3)		8	3 (37.5)	

AD, atopic dermatitis; BSA, body surface area; DLQI, Dermatology Life Quality Index; EASI, Eczema Area and Severity Index; SCORAD, Scoring Atopic Dermatitis; NRS, numeric rating scale; IGA, Investigator's Global Assessment

### Tralokinumab response in dupilumab-experienced patients

- Among dupilumab-experienced patients at Week 16, 100% (n/N, 6/6) of patients receiving tralokinumab + TCS achieved EASI-75 without the use of rescue therapy, compared to 50% (4/8) of those receiving placebo + TCS (difference [95% CI]: 50.0 [15.4, 84.6]; Table 2)
- Numerically higher proportions of dupilumab-experienced patients receiving tralokinumab + TCS achieved IGA 0/1 (4/6, 66.7%; placebo + TCS: 3/8, 37.5%; difference: 29.2 [-21.3, 79.6]) and improvement in worst daily pruritus NRS (weekly average) ≥4 points (3/6, 50%; placebo + TCS: 3/8, 37.5%; difference: 12.5 [-39.7, 64.7]) at Week 16
- Similarly, at Week 26, numerically higher proportions of dupilumab-experienced patients receiving tralokinumab + TCS achieved EASI-75 (6/6, 100%; placebo + TCS: 3/8, 37.5%; difference: 62.5 [29.0, 96.0]), IGA 0/1 (4/6, 66.7%; placebo + TCS: 2/8, 25%; difference: 41.7 [-6.5, 89.9]), and improvement in worst daily pruritus NRS (weekly average) ≥4 points (3/6, 50%; placebo + TCS: 3/8, 37.5%; difference: 12.5 [-39.7, 64.7]), compared to placebo + TCS

Table 2. Binary efficacy endpoints in dupilumab-experienced patients.

Visit	Endpoint	Tralokinumab + TCS (n=6)	Placebo + TCS (n=8)	Difference (95% CI) <sup>†</sup>
Week 16	EASI75	6 / 6 (100.0%)	4 / 8 (50.0%)	50.0 (15.4, 84.6)
	IGA 0/1	4 / 6 (66.7%)	3 / 8 (37.5%)	29.2 (-21.3, 79.6)
	Itch NRS ≥4*	3 / 6 (50.0%)	3 / 8 (37.5%)	12.5 (-39.7, 64.7)
Week 26	EASI75	6 / 6 (100.0%)	3 / 8 (37.5%)	62.5 (29.0, 96.0)
	IGA 0/1	4 / 6 (66.7%)	2 / 8 (25.0%)	41.7 (-6.5, 89.9)
	Itch NRS ≥4*	3 / 6 (50.0%)	3 / 8 (37.5%)	12.5 (-39.7, 64.7)

<sup>†</sup>Estimated treatment difference and 95% CI from Mantel-Haenszel analysis with treatment as only strata  
\*Improvement in worst daily pruritus NRS (weekly average) ≥4 points from baseline

## Safety

- Through the 26 weeks, 66.7% (4/6) of dupilumab-experienced patients receiving tralokinumab + TCS reported any adverse event, compared to 87.5% (7/8) of those receiving placebo + TCS (Table 3)
- One patient who received placebo reported 2 events of conjunctivitis, 1 mild and 1 of moderate severity; 1 tralokinumab patient reported 1 mild event of conjunctivitis
- No serious adverse events occurred in either treatment group
- From a safety perspective, there were 2 patients who had previously discontinued dupilumab due to conjunctivitis; adverse events of conjunctivitis were not reported for either patient during 26 weeks of tralokinumab + TCS treatment

Table 3. Adverse events in dupilumab-experienced patients through 26 weeks

Adverse event, n (%)	Tralokinumab + TCS (n=6)	Placebo + TCS (n=8)
Any adverse event	4/6 (66.7)	7/8 (87.5)
Any serious adverse event	0/6 (0.0)	0/8 (0.0)
Conjunctivitis*	1/6 (16.7)	1/8 (12.5)

\*Search was done for adverse event of special interest: conjunctivitis

## Conclusions

This post hoc subgroup analysis indicates that dupilumab-experienced patients can benefit from tralokinumab + TCS as needed.

- Overall frequencies of adverse events in dupilumab-experienced patients treated with tralokinumab + TCS as needed were consistent with those in the pooled analysis of tralokinumab Phase 2 and 3 trials<sup>8</sup>
- Due to the small sample size, further data involving more patients are needed to confirm these findings

## References

- Weidinger S, Novak N. *Lancet*. 2016;387:1109–22;
- Eckert L, et al. *J Am Acad Dermatol*. 2017;77:274–9e273;
- Silverberg JI, et al. *Ann Allergy Asthma Immunol*. 2018;121:340–7;
- Dalgard FJ, et al. *J Invest Dermatol*. 2015;135:984–91;
- Bieber T. *Allergy*. 2020;75:54–62;
- Tsoi LC, et al. *J Invest Dermatol*. 2019;139:1480–9;
- Popovic B, et al. *J Mol Biol*. 2017;429:208–19;
- Simpson et al. Safety of specifically targeting interleukin 13 with tralokinumab in adult patients with moderate-to-severe atopic dermatitis: pooled analysis of five randomised, double-blind, placebo-controlled Phase 3 and Phase 2 trials. Poster presented at: 29th EADV Congress; 29–31 October 2020; Virtual.

## Disclosures

Jan Gutermuth reports honoraria as a consultant/advisory board member/ speaker and/or received grants from AbbVie, Genzyme, LEO Pharma, Lilly, Pfizer, Regeneron, and Sanofi. Andrew E. Pink has acted as an advisor/speaker for AbbVie, Almirall, Janssen, La Roche-Posay, LEO Pharma, Lilly, Novartis, Pfizer, Sanofi, and UCB. Margitta Worm has served as a scientific advisor and/or clinical trial investigator and/or paid speaker for AbbVie, ALK, Allergopharma, Aimmune, Boehringer Ingelheim, Dermira, Eli Lilly, Galderma, Janssen, LEO Pharma, Mylan, Novartis, Pfizer, Regeneron and Sanofi-Genzyme. Lise Soldbro, Thomas Mark, Joshua Corriveau, Christian Bjerregård Øland, and Shannon Schneider are employees of LEO Pharma. Stephan Weidinger is co-principal investigator of the German Atopic Eczema Registry TREATgermany. He has received institutional research grants from Sanofi Deutschland GmbH, LEO Pharma, and La Roche-Posay, has performed consultancies for Sanofi-Genzyme, Regeneron, LEO Pharma, AbbVie, Pfizer, Eli Lilly, Kymab and Novartis, he has also lectured at educational events sponsored by Sanofi-Genzyme, Regeneron, LEO Pharma, AbbVie, Novartis and Galderma, and is involved in performing clinical trials with many pharmaceutical industries that manufacture drugs used for the treatment of psoriasis and atopic eczema.

**Acknowledgments** The ECZTRA 7 clinical trial was sponsored by LEO Pharma A/S, Ballerup, Denmark. Originally presented at Revolutionizing Atopic Dermatitis (RAD), June 13, 2021